

TWO NEW ANALYTICAL SOLUTIONS AND TWO NEW GEOMETRICAL SOLUTIONS FOR THE WEIGHTED FERMAT-TORRICELLI PROBLEM IN THE EUCLIDEAN PLANE

ANASTASIOS N. ZACHOS

ABSTRACT. We obtain two analytic solutions for the weighted Fermat-Torricelli problem in the Euclidean Plane which states that: Given three points in the Euclidean plane and a positive real number (weight) which correspond to each point, find the point such that the sum of the weighted distances to these three points is minimized. Furthermore, we give two new geometrical solutions for the the weighted Fermat-Torricelli problem (weighted Fermat-Torricelli point), by using the floating equilibrium condition of the weighted Fermat-Torricelli problem (first geometric solution) and a generalization of Hofmann's rotation proof under the condition of equality of two given weights (second geometric solution).

1. INTRODUCTION

We state the weighted Fermat-Torricelli problem in \mathbb{R}^2 :

Problem 1. *Given a triangle $\triangle A_1A_2A_3$ with vertices $A_1 = (x_1, y_1)$, $A_2 = (x_2, y_2)$, $A_3 = (x_3, y_3)$, find a fourth point $A_F = (x_F, y_F)$ which minimizes the objective function*

$$f(x, y) = \sum_{i=1}^3 B_i \sqrt{(x - x_i)^2 + (y - y_i)^2} \quad (1.1)$$

where B_i is a positive real number (weight) which corresponds to A_i .

By replacing $B_1 = B_2 = B_3$ in (1.1), we obtain the (unweighted) Fermat-Torricelli problem which was first stated by Pierre de Fermat (1643).

The solution of the weighted Fermat-Torricelli problem (Problem 1) is called the weighted Fermat-Torricelli point A_F .

1991 *Mathematics Subject Classification.* 51M04, 51E10.

Key words and phrases. weighted Fermat-Torricelli problem, weighted Fermat-Torricelli point.

The existence and uniqueness of the weighted Fermat-Torricelli point and a complete characterization of the "floating case" and "absorbed case" has been established by Y. S Kupitz and H. Martini (see [9], theorem 1.1, reformulation 1.2 page 58, theorem 8.5 page 76, 77). A particular case of this result for three non-collinear points in \mathbb{R}^2 , is given by the following theorem:

Theorem 1. [1],[9] *Let there be given a triangle $\triangle A_1A_2A_3$, $A_1, A_2, A_3 \in \mathbb{R}^2$ with corresponding positive weights B_1, B_2, B_3 .*

- (a) *The weighted Fermat-Torricelli point A_F exists and is unique.*
 (b) *If for each point $A_i \in \{A_1, A_2, A_3\}$*

$$\left\| \sum_{j=1, j \neq i}^3 B_j \vec{u}(A_i, A_j) \right\| > B_i, \quad (1.2)$$

for $i, j = 1, 2, 3$ holds, then

- (b₁) *the weighted Fermat-Torricelli point A_F (weighted floating equilibrium point) does not belong to $\{A_1, A_2, A_3\}$ and*
 (b₂)

$$\sum_{i=1}^3 B_i \vec{u}(A_F, A_i) = \vec{0}, \quad (1.3)$$

where $\vec{u}(A_k, A_l)$ is the unit vector from A_k to A_l , for $k, l \in \{0, 1, 2, 3\}$ (Weighted Floating Case).

- (c) *If there is a point $A_i \in \{A_1, A_2, A_3\}$ satisfying*

$$\left\| \sum_{j=1, j \neq i}^3 B_j \vec{u}(A_i, A_j) \right\| \leq B_i, \quad (1.4)$$

then the weighted Fermat-Torricelli point A_F (weighted absorbed point) coincides with the point A_i (Weighted Absorbed Case).

A direct consequence of the weighted floating case and the weighted absorbed case of theorem 1 gives Corollary 1 (Torricelli's theorem) and Corollary 1 (Cavalieri's alternative) (see in [1], p 236).

Corollary 1. *If $B_1 = B_2 = B_3$ and all three angles of the triangle $\triangle A_1A_2A_3$ are less than 120° , then A_F is the isogonal point (interior point) of $\triangle A_1A_2A_3$ whose sight angle to every side of $A_1A_2A_3$ is 120° (Torricelli's theorem).*

Corollary 2. *If $B_1 = B_2 = B_3$ and one of the angles of the triangle $\triangle A_1A_2A_3$ is equal or greater than 120° , then A_F is the vertex of the obtuse angle of $\triangle A_1A_2A_3$ (Cavalieri's alternative).*

Concerning the solution of the weighted Fermat-Torricelli problem with the use of analytic geometry and trigonometry, we mention the works of [3], [8], [10] [5], [4], [14] and [18].

Recently, an analytic solution, which express explicitly the coordinates of the weighted Fermat-Torricelli point with respect to the coordinates of the three points A_i and the three weights B_1, B_2, B_3 for the weighted Fermat-Torricelli problem with respect to the weighted floating case of Theorem 1 has been derived in [13] and for the case $B_1 = B_2 = B_3 = 1$ has been derived in [11].

In this paper, we present two new analytical solutions for the weighted Fermat-Torricelli problem in \mathbb{R}^2 in the weighted floating case of Theorem 1. The first analytical solution gives the coordinates of the weighted Fermat-Torricelli point as a function of the coordinates of the three non collinear points and the three given weights (real positive numbers) in a different way from [13] and [11] (Theorem 2, Section 2).

The second analytical solution gives the location of the weighted Fermat-Torricelli point as a function of two inscribed angles of the circumscribed circle which passes from the three non collinear points and the three given weights by applying a coordinate independent approach given in [18] (Theorem 3, Section 3).

The first geometrical solution of the weighted Fermat-Torricelli point with ruler and compass focuses on constructing the intersection of two Simpson lines (weighted case) by applying the duality of the weighted Fermat-Torricelli problem which was introduced in [13] (Problem 2, Section 4).

Finally, the second geometric solution of the weighted Fermat-Torricelli point focuses on finding the angle of rotation of the three non collinear points about one of them and generalizes Hofmann's rotation proof ([1], [6], [12]) regarding the equality of the given weights (Problem 3, Corollary 3, Section 4).

2. ANALYTICAL SOLUTION OF THE WEIGHTED FERMAT-TORRICELLI PROBLEM

We obtain an analytic solution for the floating case of Theorem 1, i.e the weighted Fermat-Torricelli point A_F is an interior point of $\triangle A_1 A_2 A_3$, such that the coordinates x_F and y_F of A_F are expressed explicitly as a function of x_i, y_i and B_i , for $i = 1, 2, 3$, by using analytic geometry in \mathbb{R}^2 .

We denote by a_{ij} the length of the linear segment $A_i A_j$ and α_{ikj} the angle $\angle A_i A_k A_j$ for $i, j, k = 0, 1, 2, 2', 3, 3', i \neq j \neq k$ (See fig. 1).

Without loss of generality, we set $A_1 = (0, 0)$, $A_2 = (a_{12}, 0)$, $A_3 = (x_3, y_3)$.

We need the following two lemmata:

Lemma 1. [4],[14] *Under the condition (1.2) and the weighted floating equilibrium condition (1.3) the following equation is satisfied:*

$$\frac{B_i}{\sin \alpha_{203}} = \frac{B_2}{\sin \alpha_{103}} = \frac{B_3}{\sin \alpha_{102}} = C, \quad (2.1)$$

$$\text{where } C = \frac{2B_1B_2B_3}{\sqrt{(B_1+B_2+B_3)(B_2+B_3-B_1)(B_1+B_3-B_2)(B_1+B_2-B_3)}}$$

Lemma 2. [1],[5],[4],[14] *Under the condition (1.2) and the weighted floating equilibrium condition (1.3) the angle α_{i0j} is expressed as a function of B_1, B_2 and B_3 :*

$$\alpha_{i0j} = \arccos \left(\frac{B_k^2 - B_i^2 - B_j^2}{2B_iB_j} \right) \quad (2.2)$$

for $i, j, k = 1, 2, 3$, and $i \neq j \neq k$.

Theorem 2. *Under the condition (1.2) of the weighted floating case, the coordinates of the weighted Fermat-Torricelli point $A_F (x_F, y_F)$ are given by the following relations:*

$$x_F = -\frac{(a_{12} - x'_2)(x'_3y_3 - x_3y'_3) + d_3(x_3 - x'_3)}{(a_{12} - x'_2)(y'_3 - y_3) - (y_3 - x'_3)y'_2}, \quad (2.3)$$

$$y_F = \frac{y'_2(a_{12}y_3 - x'_3y_3 - a_{12}y'_3 + x_3y'_3)}{x_3y'_2 - x_3y'_2 + a_{12}y_3 - x'_2y_3 - a_{12}y'_3 + x'_2y'_3} \quad (2.4)$$

where

$$x'_2 = -\frac{B_3}{B_2} \left(x_3 \frac{B_1^2 - B_2^2 - B_3^2}{2B_2B_3} + y_3 \sqrt{1 - \left(\frac{B_1^2 - B_2^2 - B_3^2}{2B_2B_3} \right)^2} \right), \quad (2.5)$$

$$y'_2 = \frac{B_3}{B_2} \left(x_3 \sqrt{1 - \left(\frac{B_1^2 - B_2^2 - B_3^2}{2B_2B_3} \right)^2} - y_3 \frac{B_1^2 - B_2^2 - B_3^2}{2B_2B_3} \right), \quad (2.6)$$

$$x'_3 = -a_{12} \frac{B_2}{B_3} \left(\frac{B_1^2 - B_2^2 - B_3^2}{2B_2B_3} \right) \quad (2.7)$$

and

$$y'_3 = -a_{12} \frac{B_2}{B_3} \left(\sqrt{1 - \left(\frac{B_1^2 - B_2^2 - B_3^2}{2B_2B_3} \right)^2} \right). \quad (2.8)$$




FIGURE 1.

Proof of Theorem 2: We apply the weighted Torricelli configuration which is similar to the configuration used in [2] and we construct two similar triangles $\triangle A_1A_2A_{3'}$ and $\triangle A_1A_3A_{2'}$, such that:

$$\alpha_{13'2} = \alpha_{132'} = \pi - \alpha_{102}, \quad (2.9)$$

$$\alpha_{213'} = \alpha_{312'} = \pi - \alpha_{203}, \quad (2.10)$$

and

$$\alpha_{123'} = \alpha_{12'3} = \pi - \alpha_{103}. \quad (2.11)$$

From (2.11), (2.11) and (2.11), we derive that the point of intersection of the two circles which pass from $A_1, A_{3'}, A_2$ and $A_1, A_{2'}, A_3$, respectively, is the weighted Fermat-Torricelli point A_F (fig. 1). Therefore, we obtain that A_F is the intersection point of the lines (weighted Simpson lines) defined by $A_2A_{2'}$ and $A_3A_{3'}$.

Thus, we have:

$$x'_2 = a_{12'} \cos(\alpha_{213} + \pi - \alpha_{203}), \quad (2.12)$$

$$y'_2 = a_{12'} \sin(\alpha_{213} + \pi - \alpha_{203}), \quad (2.13)$$

$$x'_3 = a_{13'} \cos(\pi - \alpha_{203}), \quad (2.14)$$

and

$$y'_3 = -a_{13'} \sin(\pi - \alpha_{203}). \quad (2.15)$$

By applying the sine law in $\triangle A_1 A_2 A_{3'}$ and $\triangle A_1 A_3 A_{2'}$, we get, respectively,:

$$a_{13'} = a_{12} \frac{B_2}{B_3}. \quad (2.16)$$

and

$$a_{12'} = a_{13} \frac{B_3}{B_2}. \quad (2.17)$$

By replacing (2.16), (2.17) and (2.2) from lemma 2 in (2.12), (2.13), (2.14) and (2.15), we obtain (2.5), (2.6), (2.7) and (2.8).

The equations of the lines defined by $A_2 A_{2'}$ and $A_3 A_{3'}$, respectively, are as follows:

$$\frac{y'_2}{x'_2 - a_{12}} = \frac{y}{x - a_{12}} \quad (2.18)$$

and

$$\frac{y_3 - y'_3}{x_3 - x'_3} = \frac{y_3 - y}{x_3 - x} \quad (2.19)$$

Solving (2.18) and (2.19) with respect to (x, y) we derive the point of intersection $A_F = (x_F, y_F)$, and the coordinates x_F and y_F are given by (2.3) and (2.4), respectively.

□

3. AN EXPLICIT ANGULAR SOLUTION OF THE WEIGHTED FERMAT-TORRICELLI PROBLEM

It is well known that the barycenter A_m of $\triangle A_1 A_2 A_3$ is constructed by the relation $a_{im} = \frac{2}{3}a_{i,jk} = \frac{1}{3}\sqrt{2a_{ij}^2 + 2a_{ik}^2 - a_{jk}^2}$, where $a_{i,jk}$ is the length of the midline that connects the vertex A_i with the midpoint of the line segment $A_j A_k$ for $i, j, k = 1, 2, 3$, $i \neq j \neq k$ and the median minimizes the objective function $a_{m1}^2 + a_{m2}^2 + a_{m3}^2$.

A natural question to ask is if the location of the weighted Fermat-Torricelli problem could be given with respect to the lengths of the sides of $\triangle A_1 A_2 A_3$ and the constant positive weights B_1, B_2, B_3 .

A positive answer to this question is given by the following lemma ([18, Corollary 2]):

Lemma 3. ([18, Corollary 2]) *The explicit solution of the weighted Fermat-Torricelli problem in \mathbb{R}^2 , under the condition (1.2) (weighted floating case) is given by:*

$$\alpha_{013} = \operatorname{arccot} \left(\frac{\sin(\alpha_{213}) - \cos(\alpha_{213}) \cot(\arccos \frac{B_3^2 - B_1^2 - B_2^2}{2B_1B_2}) - \frac{a_{13}}{a_{12}} \cot(\arccos \frac{B_2^2 - B_1^2 - B_3^2}{2B_1B_3})}{-\cos(\alpha_{213}) - \sin(\alpha_{213}) \cot(\arccos \frac{B_3^2 - B_1^2 - B_2^2}{2B_1B_2}) + \frac{a_{13}}{a_{12}}} \right) \quad (3.1)$$

and

$$a_{10} = \frac{\sin \left(\alpha_{013} + \arccos \frac{B_2^2 - B_1^2 - B_3^2}{2B_1B_3} \right) a_{13}}{\sin \left(\arccos \frac{B_2^2 - B_1^2 - B_3^2}{2B_1B_3} \right)}. \quad (3.2)$$

where

$$\alpha_{213} = \arccos \left(\frac{a_{12}^2 + a_{13}^2 - a_{23}^2}{2a_{12}a_{13}} \right) \quad (3.3)$$

and α_{013} and a_{10} depend on $B_1, B_2, B_3, a_{13}, a_{12}$ and a_{23} .

Proof of Lemma 3: We assume that the weighted floating case occurs (see theorem 1, Case b), in order to locate it in the interior of the $\triangle A_1A_2A_3$.

From the cosine law in $\triangle A_1A_0A_2$, and $\triangle A_1A_0A_3$ we get, respectively:

$$a_{02}^2 = a_{01}^2 + a_{12}^2 - 2a_{01}a_{12} \cos(\alpha_{213} - \alpha_{013}) \quad (3.4)$$

and

$$a_{03}^2 = a_{01}^2 + a_{13}^2 - 2a_{01}a_{13} \cos(\alpha_{013}). \quad (3.5)$$

From (3.4) and (3.5), a_{02} and a_{03} are expressed with respect to the two variables a_{01} and α_{013} :

$$a_{0i} = a_{0i}(a_1, \alpha_{013}),$$

for $i = 2, 3$. By differentiating (1.1) with respect to a_{01} and α_{013} , respectively, we get:

$$B_1 + B_2 \frac{\partial a_{02}}{\partial a_{01}} + B_3 \frac{\partial a_{03}}{\partial a_{01}} = 0, \quad (3.6)$$

$$B_2 \frac{\partial a_{02}}{\partial \alpha_{013}} + B_3 \frac{\partial a_{03}}{\partial \alpha_{013}} = 0. \quad (3.7)$$

From Appendix A, by replacing (A.1) and (A.2) in (3.6), we obtain:

$$B_2 \cos(\alpha_{102}) + B_3 \cos(\alpha_{103}) = -B_1 \quad (3.8)$$

By replacing (A.5) and (A.6) in (3.7), we obtain:

$$-B_2 \sin(\alpha_{102}) + B_3 \sin(\alpha_{103}) = 0 \quad (3.9)$$

By squaring both parts of (3.8) and (3.9) and by adding the two derived equations, we get:

$$\cos(\alpha_{203}) = \frac{B_1^2 - B_2^2 - B_3^2}{2B_2B_3}. \quad (3.10)$$

Similarly by expressing the objective function with respect to the two variables a_2, α_{023} , and with respect to the two variables a_3, α_{031} , we derive, respectively:

$$\cos(\alpha_{103}) = \frac{B_2^2 - B_1^2 - B_3^2}{2B_1B_3} \quad (3.11)$$

and

$$\cos(\alpha_{102}) = \frac{B_3^2 - B_1^2 - B_2^2}{2B_1B_2}. \quad (3.12)$$

From the sine law in $\triangle A_1A_0A_2$, $\triangle A_1A_0A_3$, we get, respectively:

$$\frac{a_{12}}{\sin(\alpha_{102})} = \frac{a_{01}}{\sin(\alpha_{213} - \alpha_{013} + \alpha_{102})} \quad (3.13)$$

and

$$\frac{a_{13}}{\sin(\alpha_{103})} = \frac{a_{01}}{\sin(\alpha_{013} + \alpha_{103})} \quad (3.14)$$

By eliminating a_{01} from (3.13) and (3.14), we obtain:

$$\alpha_{013} = \arccot\left(\frac{\sin(\alpha_{213}) - \cos(\alpha_{213}) \cot(\alpha_{102}) - \frac{a_{31}}{a_{12}} \cot(\alpha_{103})}{-\cos(\alpha_{213}) - \sin(\alpha_{213}) \cot(\alpha_{102}) + \frac{a_{31}}{a_{12}}}\right) \quad (3.15)$$

By replacing (3.12) and (3.11) in (3.15), we obtain (3.1). From the sine law in $\triangle A_1A_0A_3$, we derive (3.2).

The values of a_{01} and α_{013} give the location of the weighted Fermat-Torricelli point A_F . \square

Remark 1. *The explicit solution of the weighted Fermat-Torricelli problem is similar with the definition of a complex number in a polar form:*

$z = r \exp(i(\alpha_{213} - \alpha_{013}))$, where the absolute value of z is $r = a_1$ and the argument of z is $\arg z = \alpha_{213} - \alpha_{013}$.

Let $C(Q, R)$ be the inscribed circle with center Q and radius R which passes from the vertex A_i , for $i = 1, 2, 3$.

Each of the three central angles is given by the relation:

$$c_{iQj} = 2\alpha_{imj}, \quad (3.16)$$

such that:

$$c_{1Q2} + c_{2Q3} + c_{1Q3} = 2\pi$$

or

$$c_{1Q2} = 2\pi - c_{1Q3} - c_{2Q3}, \quad (3.17)$$

for $i \neq m \neq j$, $i, m, j = 1, 2, 3$. From the sine law in $\triangle A_1 A_2 A_3$ and taking into account (3.16), we get:

$$\frac{a_{13}}{\sin(\frac{c_{1Q3}}{2})} = \frac{a_{12}}{\sin(\frac{c_{1Q2}}{2})} = 2R. \quad (3.18)$$

By replacing (3.16), (3.17) (3.18) in (3.1) and (3.2) of lemma 3, we derive the following result:

Theorem 3. *An explicit angular solution of the weighted Fermat-Torricelli problem in \mathbb{R}^2 , under the condition (1.2) is given by:*

$$\cot \alpha_{013} = \frac{\sin(\frac{c_{2Q3}}{2}) - \cos(\frac{c_{2Q3}}{2}) \cot(\arccos \frac{B_3^2 - B_1^2 - B_2^2}{2B_1 B_2}) - \frac{\sin(\frac{c_{1Q3}}{2})}{\sin(\frac{c_{1Q3} + c_{2Q3}}{2})} \cot(\arccos \frac{B_2^2 - B_1^2 - B_3^2}{2B_1 B_3})}{-\cos(\frac{c_{2Q3}}{2}) - \sin(\frac{c_{2Q3}}{2}) \cot(\arccos \frac{B_3^2 - B_1^2 - B_2^2}{2B_1 B_2}) + \frac{\sin(\frac{c_{1Q3}}{2})}{\sin(\frac{c_{1Q3} + c_{2Q3}}{2})}} \quad (3.19)$$

and

$$a_{10} = 2R \frac{\sin(\alpha_{013} + \arccos \frac{B_2^2 - B_1^2 - B_3^2}{2B_1 B_3}) \sin(\frac{c_{1Q3}}{2})}{\sin(\arccos \frac{B_2^2 - B_1^2 - B_3^2}{2B_1 B_3})}, \quad (3.20)$$

where α_{013} and a_{10} depend on $B_1, B_2, B_3, c_{1Q3}, c_{2Q3}$, and R .

Remark 2. *We conclude that by setting $R = 1$ (unit radius circumscribed circle) in (3.20), the explicit solution depends only on five given elements: B_1, B_2, B_3, c_{1Q3} and c_{2Q3} . This unique result holds only if the inequalities (1.2) of the weighted floating case of Theorem 1, Case (b) are satisfied.*


Remark 3. *We note that lemma 3 and theorem 3 provide an analytic solution for the weighted Fermat-Torricelli problem in \mathbb{R}^2 without using the coordinates of the points A_i , for $i = 1, 2, 3$ (Coordinate independent approach) taking into account only given Euclidean elements (lengths and angles).*

4. TWO NEW GEOMETRICAL SOLUTIONS OF THE WEIGHTED FERMAT-TORRICELLI POINT IN THE EUCLIDEAN PLANE

We present two new geometrical solutions to find the weighted Fermat-Torricelli point in the weighted floating case.

The first solution deals with the position of $A_{1'}$ and $A_{3'}$ which shall give the position of the two Simpson lines defined by A_2 and $A_{2'}$ and $A_3, A_{3'}$ (fig. 2)

The second solution deals with the generalization of Hofmann's rotation proof ([12],[6]) for $B_1 = B_2$, such that B_1



Problem 2. *Construct the solution of Problem 1 (Weighted Fermat-Torricelli problem) using ruler and compass, under the condition (1.2).*

Take a point G to the line defined by A_1A_2 , such that $\|A_1A_2\| = a_{12}$, $\|A_2G\| = B_3$ where $\|A_1G\| = a_{12} + B_3$ and construct with a ruler and compass the perpendicular linear segment at the point G to the line defined by A_1A_2 and take a point H such that $\|GH\| = B_2$. We denote by I the point of intersection of the line defined by A_2H and the perpendicular line at the point A_1 with respect to the line defined by A_1A_2 (fig. 3). Taking into account the similar triangles $\triangle A_1A_2I$ and $\triangle A_2GH$, we get:

Similarly, we construct the vertex $A_{2'}$. We select a point M which belongs to the linear segment A_1A_3 , such that $\|A_1M\| = B_2$ and we




FIGURE 3.

construct a triangle $\triangle A_1MN$ with the other two sides $\|A_1N\| = B_3$ and $\|MN\| = B_1$.

Thus, lemma 1 yields $\angle A_3A_1A_{2'} = \alpha_{312'} = \pi - \alpha_{203}$.

We need to calculate $a_{12'}$, in order to find the location of $A'_{2'}$.

Take a point Q to the line defined by A_1A_3 , such that $\|A_1A_3\| = a_{13}$, $\|A_2Q\| = B_3$ where $\|A_1Q\| = a_{13} + B_3$ and construct with a ruler and compass the perpendicular linear segment at the point Q to the line defined by A_1A_3 and take a point R such that $\|RQ\| = B_3$. We denote by P the point of intersection of the line defined by A_3R and the perpendicular line at the point A_1 with respect to the line defined by A_1A_3 (fig. 4). Taking into account the similar triangle $\triangle A_1A_3P$ and $\triangle A_3QR$, we get:

$$a_{12'} = a_{13} \frac{B_3}{B_2}. \quad (4.2)$$

□

Problem 3. Solve Problem 1 (Weighted Fermat-Torricelli problem) by generalizing Hofmann's rotation, under the condition (1.2) and $B_1 = B_2$, such that $B_1 + B_2 + B_3 = 1$, $B_1 > \frac{1}{4}$ and $\alpha_{132} > \pi - \arccos\left(-1 + \frac{(1-2B_1)^2}{2B_1^2}\right)$.

Solution of Problem 3: We consider a weight B_i which corresponds to the vertex A_i in \mathbb{R}^2 , for $i = 1, 2, 3$.

By replacing $B_1 = B_2$ in (2.2) of lemma 2, we derive that:




FIGURE 4.




FIGURE 5.

$$\alpha_{203} = \alpha_{103} = \arccos \left(1 - \frac{1}{2B_1} \right) \quad (4.3)$$

and

$$\alpha_{102} = \arccos \left(-1 + \frac{(1 - 2B_1)^2}{2B_1^2} \right) \quad (4.4)$$

which yields $B_1 > \frac{1}{4}$.

Taking into account (4.3) and (4.4), we rotate the triangle $\triangle A_1 A_2 A_3$ about A_3 through $\pi - \alpha_{102} = 2\alpha_{103} - \pi \text{ rad}$ and we obtain the triangle $\triangle A_3 A'_1 A'_2$. Let A'_F be the corresponding weighted Fermat-Torricelli point of $\triangle A'_1 A'_2 A_3$, for $B'_1 = B_1$ and $B_2 = B'_2$ (fig. 5). Thus, the points A_2 , A_F , A'_F and A'_3 are collinear (fig. 5), because $\triangle A_F A'_F A_3$ is an isosceles triangle and

$$\angle A_F A'_F A_3 = \angle A'_F A_F A_3 = \pi - \alpha_{103}.$$

□

Corollary 3. [6], [12], [1] *For $B_1 = B_2 = B_3$ the solution of Problem 3 is given by rotating the $\triangle A_1 A_2 A_3$ about A_3 through 60° .*

Proof. By replacing $B_1 = B_2 = B_3 = 1$ in the solution of Problem 3, we deduce that the rotation about A_3 need to be $\pi - 120^\circ = 60^\circ < \alpha_{132}$ (Hofmann's rotation). □

The author is sincerely grateful to Professor Dr. Vassilios G. Papa-georgiou for his very valuable comments, fruitful discussions and for his permanent attention to this work.

APPENDIX A.

We mention two methods of the length of a linear segment with respect to (I) a variable length and (II) a variable angle, which have been used, in order to find the weighted Fermat-Torricelli point.

I. A method of differentiating the length of a linear segment with respect to the length of a variable linear segment is given first in [14, Proposition 2.6, (b), Remark 2.4, Corollary 3.3], [15, formula (4), p. 413] and has been explained in detail in [17], [16, Corollary 2]. Specifically, by differentiating (3.4) with respect to a_1 , and by replacing in the derived equation $\cos(\alpha_{213} - \alpha_{013})$ taken from (3.4), we obtain:

$$\frac{\partial a_2}{\partial a_1} = \cos(\alpha_{102}). \quad (\text{A.1})$$

Similarly, by differentiating (3.5) with respect to a_1 , and by replacing in the derived equation $\cos(\alpha_{013})$ taken from (3.5), we obtain:

$$\frac{\partial a_3}{\partial a_1} = \cos(\alpha_{103}). \quad (\text{A.2})$$

We mention a method of differentiating the length of a linear segment with respect to a variable angle, which have been used, in order to find the weighted Fermat-Torricelli point ([14, Proposition 2.6 (b)]) in \mathbb{R}^2 . By mentioning this technique of differentiation, we correct some typographical errors which appear in [14]. Specifically, by differentiating (3.4) with respect to α_{013} , we get:

$$\frac{\partial a_{02}}{\partial \alpha_{013}} = -a_{01} \frac{a_{12}}{a_{02}} \sin(\alpha_{213} - \alpha_{013}) \quad (\text{A.3})$$

From the sine law in $\triangle A_1 A_0 A_2$, we get:

$$\frac{a_{12}}{\sin(\alpha_{102})} = \frac{a_{02}}{\sin(\alpha_{213} - \alpha_{013})} \quad (\text{A.4})$$

By replacing (A.4) in (A.3), we obtain:

$$\frac{\partial a_{02}}{\partial \alpha_{013}} = -a_{01} \sin(\alpha_{102}). \quad (\text{A.5})$$

Similarly, by differentiating (3.5) with respect to α_{013} , we obtain:

$$\frac{\partial a_{03}}{\partial \alpha_{013}} = a_{01} \sin(\alpha_{103}). \quad (\text{A.6})$$

REFERENCES

- [1] V. Boltyanski, H. Martini, V. Soltan, *Geometric Methods and Optimization Problems*, Kluwer, Dordrecht-Boston-London, 1999.
- [2] E.Engelbrecht, *Planimetrischer Lehrsatz*. Arch. Math. Phys. **60**(1877), 447–448.
- [3] I. Greenberg and R. Robertello, *The Three Factory Problem*, Math. Mag. **67**, no.1 (1965), 67-72.
- [4] S. Gueron and R. Tessler, *The Fermat-Steiner problem*, Amer. Math. Monthly **109** (2002), 443-451.
- [5] M. Hajja, *An Advanced Calculus Approach to finding the Fermat Point*, Math. Mag. **67**, no.1 (1994), 29-34.
- [6] J. Hofmann, Hofmann, *Elementare Losung einer Minimumsaufgabe*. Zeitschr. f. math. u. naturw. Unterricht **60** (1929) 22-23.
- [7] R. Honsberger, *Mathematical Gems I*. The Dolciani Mathematical Expositions, No.1 Math. Asoc. America, 1973.
- [8] E. Jones, *The 3-vertex single source Weber location problem*, International Journal of Mathematical Education in Science and Technology, **19**, no.5, 671-679.

- [9] Y.S. Kupitz and H. Martini, *Geometric aspects of the generalized Fermat-Torricelli problem*, Bolyai Society Mathematical Studies.**6**, (1997) , 55-127.
- [10] G. Pesamosca, *On the analytic solution of the 3-point Weber problem*, Rend. Mat. Appl. **7**, no. 11 , no. 1 (1991), 39-45.
- [11] I. Roussos, *On the Steiner minimizing point and the corresponding algebraic system*. College Math. J. 43 (2012), no. 4, 305–308.
- [12] V.M.Tikhomirov, *Stories about maxima and minima* Translated from the 1986 Russian original by Abe Shenitzer. Mathematical World, American Mathematical Society, Providence, Mathematical Association of America, Washington DC, 1990.
- [13] A. Uteshev, *Analytical solution for the generalized Fermat-Torricelli problem*, Amer. Math. Monthly. 2014. **121**, no. 4 (2014) 318-331.
- [14] A.N. Zachos and G. Zouzoulas, *The weighted Fermat-Torricelli problem and an "inverse" problem*, J. Convex Anal, **15**, no. 1 (2008) 55–62.
- [15] A.N. Zachos and G. Zouzoulas, *An evolutionary structure of convex quadrilaterals*, J. Convex Anal., **15**, no.2 (2008) 411–426.
- [16] A. Zachos and A. Cotsiolis, *The weighted Fermat-Torricelli problem on a surface and an "inverse" problem*, J. Math. Anal. Appl., **373**, no. 1 (2011) 44–58.
- [17] A. Cotsiolis and A. Zachos, *Corrigendum to "The weighted Fermat-Torricelli problem on a surface and an "inverse" problem"*, J. Math. Anal. Appl., **376**, no. 2 (2011) 760.
- [18] A. N. Zachos, *Exact location of the weighted Fermat-Torricelli point for flat surfaces of revolution*, Results. Math. **65**, no. 1-2 (2014) 167–179.

UNIVERSITY OF PATRAS, DEPARTMENT OF MATHEMATICS, GR-26500 RION,
GREECE

E-mail address: azachos@gmail.com